

The effects of reflexology on pain and sleep deprivation in patients with rheumatoid arthritis: A randomized controlled trial

Ercan Bakir^a, Sevgin Samancioglu Baglama^{a,*}, Savas Gursoy^b

^a Gaziantep University Health Science Faculty Nursing Department, 27310, Gaziantep, Turkey

^b Gaziantep University Medical Faculty, Department Of Physical Medicine and Rehabilitation, 27310, Gaziantep, Turkey

ARTICLE INFO

Article history:

Received 12 December 2017

Received in revised form

9 February 2018

Accepted 28 February 2018

ABSTRACT

Objective: This study was intended to examine the effect of foot reflexology on RA patients' pain and sleep quality.

Methods: This is a randomized controlled trial and was held at the "Rheumatology Follow-up Polyclinic" in Turkey between January–July 2015. A total of 60 patients were included in the research. A socio-demographic data form, the Pittsburgh Sleep Quality Index (PSQI) and the Visual Analogue Scale (VAS) were used. Foot Reflexology was administered to the experimental group.

Results: The research found that the pain scores of the experimental group were statistically more significant than those of the control group ($p < .01$). The experimental group's average pain was reduced by the six weeks of foot reflexology. The total PSQI score of the experimental group was lowered.

Conclusions: Foot reflexology is a non-pharmacological nursing intervention that may reduce the pain and sleep deprivation symptoms of RA patients.

© 2018 Elsevier Ltd. All rights reserved.

1. Introduction

Rheumatoid arthritis (RA) is a chronic, idiopathic, systemic, inflammatory, autoimmune disease characterized by pain in the joints and swelling, loss of function in the joints, morning stiffness and sleep deprivation as well as fatigue accompanying these states [1]. The loss of function due to the disease also leads to poor quality of life [2]. Depending on the prognosis, physical deformities and unbearable pain may become prominent over time [3]. Pain causes poor quality sleep; furthermore, patients complain about fatigue not only because of their disease but also because of sleep deprivation [4]. The incidence of sleep disorders in individuals with RA was reported to range between 54% and 70%. Therefore, these individuals' productivity at work and skills for conducting daily practices decrease [3].

The dissatisfaction with medical treatment methods, invasive procedures and the necessity of using daily analgesics as well as the toxic and harmful effects of drugs push patients to different quests for symptom management. Non-pharmacological methods such as reflexology, massage, hydrotherapy, therapeutic touch,

acupuncture, music therapy have been used for symptom control and functional improvement in these patients [5,6].

Reflexology is the stimulation of reflex points through massage on some body parts. Foot reflexology is the practice of applying pressure to specific areas of the feet. Reflexes in the feet match to organs, glands, and systems of the body. There are some theories on the mechanism of action of reflexology [7,8]. According to the haemodynamic theory, reflexology stimulation increases blood flow to the related organ or body part. The nerve impulse theory argues that reflexology stimulation enhances nervous connection to the corresponding body parts. According to the energy theory; organs and body parts are linked through electromagnetic fields and that these pathways are blocked in states of disease. Lactic acid theory defends that lactic acid accumulates on the soles of the feet in the form of crystals and reduces regular flow. Reflexology shows that the fusion of crystals promotes free circulation [7,8]. With these features, reflexology becomes an appropriate alternative method for RA patients.

Although the effect of foot reflexology on pain symptom has been evaluated in RA patients, no study has been conducted on sleep quality. The aim of this randomized controlled trial is to analyze the effects of foot reflexology on pain and sleep quality in RA patients. Two hypotheses were explored in this study. One of the hypotheses is that foot reflexology practice reduces pain and

* Corresponding author.

E-mail addresses: e.bakir02@gmail.com (E. Bakir), sevginsamancioglu@hotmail.com (S.S. Baglama), savasgursoy@gantep.edu.tr (S. Gursoy).

the other is that foot reflexology improves sleep quality in RA patients.

2. Material and methods

2.1. Ethics

The Institutional Ethics Committee of Gaziantep University Medical Faculty (No: 2014/344) approved this study before implementation. Informed consent was obtained from all the participants included in the study.

2.2. Design and sample

This study was planned as a randomized controlled trial to evaluate the effect of foot reflexology on pain and sleep quality in RA patients. The study was held at the “Rheumatology Follow-up Polyclinic” of a university hospital located in a big city in Turkey between January–July 2015. A total of 60 patients (30 for the intervention group and 30 for the control group) who agreed to participate in the research and met the criteria specified within the population were included in the research. All the patients were suffering from pain and sleep disturbances. The patients included in the research demonstrated the following characteristics: 18 years of age or older, diagnosed with RA for at least 1 year, VAS pain score of 4 or greater and voluntary participation in the study. The exclusion criteria were as follows: acute infection or fever, vascular disease in the lower extremities, impaired skin integrity, history of surgery, fractures, sprains or injuries in the lower extremity, pregnancy, diagnosis of diabetes, diagnosis of sleep apnea and using sleep medications, cigarette and alcohol consumption. Medical treatments of RA patients in both control and reflexology groups continued routinely. However, no change was allowed in medical treatment for RA during six reflexology applications and control. An a priori power analysis was performed to calculate the sample size (STATISTICA 2016 Quest Software Inc.). As a result of power analysis, the minimum sample size required for the study was found as the effect size = .50, alpha = .05 and the power of the test (1- beta) .80, while 30 patients were included in each group. Fig. 1 shows the progression of the study (Fig. 1). Patients with similar levels of VAS pain, gender and disease duration were selected for the experimental and control groups. They were then divided into two groups as foot reflexology and control group using a random number table.

2.3. Data collection

The interviews with RA patients were conducted face to face and individually at the Rheumatology Follow-up Polyclinic. At the interviews, the sociodemographic information and biochemical parameters of the patients as well as their Visual Analogue Scale for Pain (VAS-Pain) and Pittsburgh Sleep Quality Index (PSQI) scores were collected. The form questioning socio-demographic characteristics include the age, gender, marital status, educational status, health insurance, alcohol and smoking status of the patients as well as their duration of diagnosis and the medications they take [10,11]. The pain values of both the experimental and control groups were recorded weekly for 6 weeks. The sleep quality was assessed only at the first and sixth week follow-up because the scale used in the sleep quality assessment demonstrated the last month.

VAS- Pain – Visual Analogue Scale was used to assess pain symptoms. Using a 10 cm horizontal line, a plane starting at 0 and ending at 10 points was created even intervals of 1 cm “0” means no

pain while “10” means the most severe, unbearable pain. The mean pain level during the last week was used to evaluate the baseline pain used in patient selection. Patients who had a baseline pain score of 4 or greater were admitted to the study. Current pain was assessed for weekly monitoring.

With the *Pittsburgh Sleep Quality Index*, the patients are asked 19 questions to assess their sleep quality within the past month. There are 7 components of PSQI. The assessment of each item is scored between 0 and 3 according to the frequency of symptoms. The total score is calculated including 0 and 21. Scores of six or above indicate that sleep quality is impaired [12]. The validity and reliability of the Turkish version of the scale was performed by Ağargün et al. (1996) and the Cronbach's alpha value was found to be 0.80 [13]. The Cronbach's alpha value for the current study was 0.78, indicating that the scale is highly reliable.

2.4. Interventions

Prior to any interventions and again at the completion of the interventions, all patients were asked to mark their pain level on the VAS plane. Interventions were undertaken by a certified researcher. In the experimental group, 30-min foot reflexology for a total of 6 times (once a week) was performed. After 6 weeks of reflexology, the patients in the experimental group were again subjected to VAS and PSQI assessment. All patients enrolled in the study were routinely given RA treatments. The patients, particularly in the experimental group, were reminded not to take analgesic medication during the days of intervention. As for the patients in the control group, they continued to undergo routine polyclinic monitoring which included intra-articular and extra-articular physical findings, laboratory tests, radiology examinations, and information on medication.

2.4.1. Foot reflexology protocol

The room temperature of the patients' follow-up polyclinics ranged between 71.60–75.20 ° F and the humidity ranged between 40 and 60%. For the application, colorless Vaseline with no additives was used. Beginning with the right foot, 5 min of foot warming movements were provided, 3-min massage for the brain part (application to the pituitary, pineal gland points at the big toe, from the big toe to the middle of the heel; the medulla spinalis point), 2-min pressure to the solar plexus point (the reflex point of the central nervous system on the sole), 3-min application to the lymph system, 3-min application to the diaphragm area, 3-min application to the thyroid point, 3-min application to the stomach point, 3-min application to the adrenal glands, and 5-min warming movements after the application was over which added up to a total of 30 min. Foot reflexology was performed with the same techniques on the left foot after the treatment to the right foot was completed. The entire procedure was completed with a 30 - min application for each foot which lasted for a total of 60 min. The treatment was repeated once a week for 6 weeks.

2.5. Data analysis

The data were evaluated with Statistical Package for Social Sciences 18.0 software (SPSSFW, SPSS Inc., Chicago, IL, USA SPSS). Kolmogorov Simirnov test was used to determine whether the data had a normal distribution or not, and nonparametric tests were conducted when the data did not demonstrate a normal distribution. Descriptive information on the patients was given in number, percentage and standard deviation. For the evaluation of data, Chi-Square and Significance of the Difference between Two




Fig. 1. Subject selection process and progress of the study.

Means tests were used to test the differences between the groups. For the evaluation of the analyses, $p < .05$ was considered significant.

3. Results

3.1. Socio-demographic and disease characteristics of subjects

The socio-demographic and disease characteristics of the subjects are given in Table 1. The mean age of the patients is 50.16 ± 14.32 (22–74 years). Most of the patients were female (76.6%). The majority of the patients were not well-educated (70.0%). Most of the patients were married (93.3%). Almost all of the patients had health insurance (96.7%). Some of the patients were cigarette (25%) and alcohol (6.6%) consumers. The mean duration of RA diagnosis of the patients was 12.25 ± 7.65 years. The mean erythrocyte sedimentation rate of patients was 30.12 ± 18.05 mm/h while their mean C-reactive protein (CRP) value was 27.01 ± 33.58 mg/L. Of all patients enrolled, 66.6% were being treated with steroids, 63.3% with Methotrexate (MTX), 63.3% with Sulfasalazine (SSZ) and 50% with leflunomide. The mean pain score as expressed by the patients at the first meeting was found to be 6.38 ± 1.27 (5–10) while 58.3% of the patients stated that their sleep disturbances were at a severe level.

3.2. The effect of foot reflexology on pain and sleep quality

In Table 2, the pain levels and sleep quality scores of RA patients in Foot Reflexology and control groups were compared. At the end of the 6-week period, statistically significant results were obtained regarding the VAS and PSQI scores of the foot reflexology group when compared to the control group ($p < .05$) (see Table 3).

4. Discussion

This study shows that foot reflexology applied to RA patients reduces the symptoms of pain related to the disease and improves sleep quality. In the meta-analysis study conducted by Lee et al., [2011], it was concluded that foot reflexology was a useful nursing practice that had a more positive effect on fatigue and sleep than pain [14]. In another study, it was stated that foot reflexology had a mitigating effect on the symptoms of fatigue in RA patients [15].

A limited number of previous studies have also shown that foot reflexology had a pain-reducing effect in RA patients [2,9,16]. It was found that both hand and foot reflexology had a positive effect on pain and quality of life in RA patients [2]. In a study, it was concluded that foot reflexology was a more effective method in reducing pain symptoms compared to aromatherapy [15]. It was determined that reflexology applied to RA patients for six weeks as one session per week has a positive effect on the foot pain scores

Table 1
Baseline demographic and clinical characteristics of the rheumatoid arthritis subjects (N = 60).

Characteristics	Total Sample N (%)	Foot Reflexology Group N (%)	Control Group N (%)	χ^2	<i>p</i>
Age (years)					
X ± SD	50.16 ± 14.32	50.83 ± 12.05	49.50 ± 16.42	$t = 0.358$.722
Gender					
Female	46 (76.6)	24 (80)	22 (73.3)	0.373	.542
Male	14 (24.0)	6 (20)	8 (26.7)		
Education					
5 year	42 (70.0)	21 (50.0)	21 (50.0)	1.641	.801
>5 year	18 (30.0)	9 (50.0)	9 (50.0)		
Marital Status					
Married	56 (93.3)	29 (96.7)	27 (90)	1.071	.301
Single	4 (6.7)	1 (3.3)	3 (10)		
Health insurance					
Yes	58 (96.6)	29 (96.7)	29 (96.7)	0.000	1.000
No	2 (3.4)	1 (3.3)	1 (3.3)		
ESR					
(X ± SD)	30.12 ± 18.05	25.06 ± 15.38	35.18 ± 19.32	$t = -2.245$.029
CRP					
(X ± SD)	27.01 ± 33.58	14.87 ± 15.88	39.15 ± 41.68	$t = -2.981$.004
NSAID					
Yes	53 (88.3)	28 (93.3)	25 (83.3)	1.456	.228
No	7 (11.7)	2 (6.7)	5 (16.7)		
Corticosteroids					
Yes	40 (66.6)	21 (70)	19 (63.3)	0.300	.584
No	20 (33.4)	9 (30)	11 (36.7)		
MTX					
Yes	38 (63.3)	19 (63.3)	19 (63.3)	0.000	1.000
No	22 (36.7)	11 (36.7)	11 (36.7)		
SSZ					
Yes	38 (63.3)	22 (73.3)	16 (53.3)	2.584	.108
No	22 (36.7)	8 (26.7)	14 (46.7)		
Leflunomide					
Yes	30 (50)	15 (50)	15 (50)	0.000	1.000
No	30 (50)	15 (50)	15 (50)		
Pain					
(X ± SD)	6.38 ± 1.27	5.63 ± 1.06	7.13 ± 1.00	$t = -5.599$.001
Sleep Problem					
Moderate	25 (41.6)	13 (43.3)	12 (40)	0.069	0.793
Severe	35 (58.4)	17 (56.7)	18 (60)		

ESR: erythrocyte sedimentation rate; CRP: c-reactive protein; NSAID: nonsteroidal anti-inflammatory drugs; MTX: methotrexate; SSZ: sulfasalazine.

Table 2
Comparison of pain and sleep quality characteristics of groups after the first and sixth week follow-up.

	Foot Reflexology Group (n = 30)	Control Group (n = 30)	<i>T</i>	<i>p</i>
VAS-Pain				
1st Week Follow-up	7.4 ± 1.13	8.1 ± 1.24	-2.281	.026
6 th Week Follow-up	5.4 ± 0.89	8.1 ± 1.02	-10.847	.001
<i>t</i>	$t = -1.966$	$t = -9.781$		
<i>p</i>	$p = .059$	$p = .001$		
PSQI				
1 st Week Follow-up	16.20 ± 3.70	16.75 ± 3.64	-0.386	0.703
6 th Week Follow-up	13.16 ± 3.57	19.03 ± 3.05	-6.834	.001
<i>t</i>	$t = -2.651$	$t = -6.052$		
<i>p</i>	$p = .001$	$p = .001$		

perceived by the patients [9]. When compared to the control group, it was found that there was a significant decrease in mean pain scores in the foot reflexology group.

Sleep problems are observed in RA patients, especially due to pain [17]. Among the major sleep problems observed in these patients are difficulty in falling asleep and maintaining sleep, continuous disruption in sleep at night, waking up early in the morning, dizziness throughout the day and fatigue due to sleep deprivation [10]. Our study was supported by findings from a study by Pehlivan et al. (2016) that RA patients experience sleep problems and have poor sleep quality [10].

In the literature, we could not find any other study that analyzes the effect of foot reflexology on sleep quality in patients with RA. When compared with the foot reflexology group, sleep quality in the control group was significantly impaired after 6 weeks. It could be interpreted that this condition could be related to the increase in the mean pain score of the patients in the control group.

There are some limitations to this study. According to the sample size structure of the study, the findings cannot be generalized. Second, the placebo control group was not used in this study. Third, there was no directive on the use of analgesics in the control group. Therefore, the long-term effects of foot reflexology

Table 3

Evaluation of mean pain scores in sixth week follow-up (N = 60).

		N	VAS-Pain (X ± SD)	F	p ^a
1 st Follow-up	Reflexology	30	7.40 ± 1.13	5.205	.026
	Control	30	8.10 ± 1.24		
2 nd Follow-up	Reflexology	30	7.23 ± 0.89	14.157	.001
	Control	30	8.16 ± 1.01		
3 rd Follow-up	Reflexology	30	6.93 ± 0.86	11.139	.001
	Control	30	7.80 ± 1.12		
4 th Follow-up	Reflexology	30	6.30 ± 0.83	51.251	.001
	Control	30	8.10 ± 1.09		
5 th Follow-up	Reflexology	30	5.40 ± 0.96	105.216	.001
	Control	30	7.93 ± 0.94		
6 th Follow-up	Reflexology	30	5.40 ± 0.89	117.668	.001
	Control	30	8.10 ± 1.02		

^a Anova; significance level $p < .05$.

on pain and sleep quality of the patients in this group are not known. Finally, it remains unknown whether or not the practice of reflexology affects other symptoms experienced by RA patients. Thus, it is recommended that further research be conducted in this field.

5. Conclusions

The results of this study show that foot reflexology applied to RA patients is effective in increasing sleep quality, as well as reducing pain symptoms. This study shows that foot reflexology has an effect that improves sleep quality while also supporting the results of the previous studies conducted on the effects of reflexology on pain relief.

In this study, it was shown that pain and sleep symptoms of RA patients could be reduced by foot reflexology in a short time. Furthermore, this study also shows that nurses working in RA clinics may find it convenient to learn and use foot reflexology as one of the independent functions of nursing. The crucial point here is education and experience. If nurses receive good training and gain experience in performing foot reflexology, the success of the method may increase.

The effect of foot reflexology on other symptoms in RA patients should also be investigated and RA patients should be given a more comprehensive care.

Funding statement

This study was supported by the Gaziantep University's Scientific Research Project (BAP/SBF.15.01).

Conflicts of interest

Gaziantep University's Scientific Research Project were responsible for the analysis and documentation of the study.

Acknowledgements

We thank all patients who participated in this study.

Appendix A. Supplementary data

Supplementary data related to this article can be found at <https://doi.org/10.1016/j.ctcp.2018.02.017>.

References

- [1] Y.O. Jung, A.H. Kim, Recent paradigm shifts in the diagnosis and treatment of rheumatoid arthritis, *Korean J. Intern. Med. (Engl. Ed.)* 27 (2012) 378–387.
- [2] N.M. Taha, Z.H. Ali, Effect of reflexology on pain and quality of life in a patient with rheumatoid arthritis, *Life Sci. J.* 8 (2011) 357–365.
- [3] H.S. Smith, A.R. Smith, P. Seidner, Painful rheumatoid arthritis, *Pain Physician* 14 (5) (2011) E427–E458.
- [4] R.J. Stack, M. Sahni, C.D. Mallen, K. Raza, Symptom complexes at the earliest phases of rheumatoid arthritis: a synthesis of the qualitative literature, *Arthritis Care Res.* 65 (12) (2013) 1916–1926.
- [5] L. Senolt, W. Grassi, P. Szodoray, Laboratory biomarkers or imaging in the diagnostics of rheumatoid arthritis? *BMC Med.* 12 (1) (2014) 49.
- [6] N. Vivar, R.F. Van Vollenhoven, F1000prime reports, *Advances In The Treatment of Rheumatoid Arthritis*, 6, 2014.
- [7] S. Bezgin, B. Elbasan, Serebral Palsi Ve Refleksoloji, Hacettepe Üniversitesi Fizyoterapi Seminerleri, 2016, ISBN 978-605-88879-1-6, pp. 70–74. Sayfa.
- [8] A. Özdeğir, The effect of reflexology on the quality of life with breast cancer Patients, *Compl. Ther. Clin. Pract.* 29 (2017) 122–129.
- [9] S. Khan, S. Otter, K. Springett, The effects of reflexology on foot pain and quality of life in a patients with rheumatoid arthritis, a case report, *Foot* 16 (2) (2006) 112–116.
- [10] S. Pehlivan, A. Karadakovan, Y. Pehlivan, A.M. Onat, Sleep quality and factors affecting sleep in elderly patients with rheumatoid arthritis in Turkey, *Turk. J. Med. Sci.* 46 (4) (2016) 1114–1121.
- [11] Ş.Ç. Yücel, Y. Tokem, E.A. Korhan, S. Çalbay, Investigation into sleep and quality of life in patients with rheumatoid arthritis, *Int. J. Hum. Sci.* 11 (1) (2014) 457–468.
- [12] J.E.M. McCullough, A pilot randomised controlled trial exploring the effects of antenatal reflexology on labour outcomes, *Midwifery* 55 (2017) 137–144.
- [13] M.Y. Ağargün, H. Kara, Ö. Anlar, The validity and reliability of the pittsburgh sleep quality index, *Türk Psikiyatri Derg.* 7 (1996) 107–115.
- [14] J. Lee, M. Han, Y. Chung, J. Kim, J. Choi, Effects of foot reflexology on fatigue, sleep and pain: a systematic review and meta-analysis, *J. Kor. Acad. Nurs.* 41 (6) (2011) 821–833.
- [15] S. Otter, A. Church, A. Murray, J. Lucas, N. Creasey, J. Woodhouse, R. Grant, H. Cooper, The effect of reflexology in reducing the symptoms of fatigue in people with rheumatoid arthritis: a preliminary study, *J. Alternative Compl. Med.* 16 (12) (2010) 1251–1252.
- [16] Z. Gök Metin, L. Özdemir, The effects of aromatherapy massage and reflexology on pain and fatigue in patients with rheumatoid arthritis: a randomized controlled trial, *Pain Manag. Nurs.* 17 (2) (2016) 140–149.
- [17] A.C. Goes, L.A. Reis, M.B. Silva, B.S. Kahlow, T.L. Skare, Rheumatoid arthritis and sleep quality, *Rev. Bras. Reumatol.* 57 (4) (2017) 294–298 (English Edition).